


# AQUACOPP (COPPER)

## SLOW RELEASE WATER ACTIVE TABLETS

HIGHLY EFFECTIVE AND SIMPLE TO USE, **AQUACOPP** TABLETS (**CHELATE**) RELEASE ON AVERAGE 115 MG OF COPPER PER DAY... LOW COPPER CAUSES SIGNIFICANT ECONOMIC LOSSES WHERE IT OCCURS AND CAN BE CAUSED BY LOW COPPER IN THE DIET OR INTERFERENCE WITH COPPER ABSORPTION IN THE ANIMAL DUE TO MOLYBDENUM AND / OR SULPHATES IN THE FOOD / WATER. LABOUR FREE (NO ANIMAL HANDLING, NO INJECTIONS) **AQUACOPP** (**CHELATE**) PROVIDES DAILY, ORAL SUPPLEMENTATION AT A DOSE RATE OF 2 TABLETS PER HEAD PER WEEK.



SYMPTOMS OF LOW HERD COPPER IN CATTLE INCLUDE DIARRHOEA, UNTHRIFTY APPEARANCE, ROUGH FADED HAIR COATS, POOR WEIGHT GAIN, WEAK CALVES AT BIRTH, CALVING PROBLEMS (RETAINED CLEANINGS, SLOW CALVINGS) AND INFERTILITY (REPEAT-BREEDERS, SILENT HEAT). **AQUACOPP** (**CHELATE**) OFFER A GENEROUS AND SAFE SUPPLEMENT OF COPPER...THE CORRECT DOSE AT THE RIGHT TIME.

TO USE: PLACE TWO TABLETS PER HEAD PER WEEK FOR CATTLE AND 1 TABLET PER HEAD PER WEEK FOR CATTLE UNDER 300KG INSIDE ATTACHED NET DISPENSER. PLACE DISPENSER UNDER THE WATERLINE OF THE ANIMALS DRINKING TROUGH. REPLACE WEEKLY WITH FRESH TABLETS FOR THE DURATION OF SUPPLEMENTATION. USE **AQUACOPP** (**CHELATE**) 4 - 6 WEEKS BEFORE CALVING AND FOR 10 - 12 WEEKS AFTER CALVING, ESPECIALLY IN THE 4 WEEKS BEFORE SERVICE AND THROUGHOUT THE BREEDING SEASON.

